

UPWARD AND HORIZONTAL TAU AIR-SHOWERS FROM EARTH CRUST CROWN: TRACES OF HIGHEST ENERGY NEUTRINO BEAM DUMP IN WIDEST VOLUMES

D.FARGION

Physics department, Universita' degli studi "La Sapienza",
Piazzale Aldo Moro 5, - 00185 Roma, Italy
INFN Roma, Istituto Nazionale di Fisica Nucleare, Italy

Abstract

High Energy Neutrino signals from PeVs up to GZK, $E_\nu \geq 10^{19}$ eV, energies amplified as Upward and Horizontal Tau air Showers, **UPTAUS**, and **HORTAUS**, may flash toward mountain peaks, airplanes, balloons and satellites in future Tau Air-Showering experiments. Observing from high mountains the detectable crown masses at UHE ν energies around EeVs are comparable to a few km^3 , while observing from higher balloons or satellites at orbit altitudes, at GZK energies, the Horizontal Crown effective Masses may even exceed 150 km^3 . The consequent event rate by expected GZK or Z-Showering neutrinos models must arise at large rate by present scheduled (EUSO-like) or future (Crown Array Detector) satellite in Space.

1 INTRODUCTION: UHE ν Astronomy

Rarest TeVs gamma signals are at present the most extreme and rarest trace of High Energy Astrophysics. The TeVs signals have shown new power-full Jets blazing to us from Galactic or extragalactic edges. At PeVs energies astrophysical Gamma cosmic rays should also be present, but, excluded a very rare and elusive CygX3 event, they are not longer being observed. While the corresponding PeVs charged cosmic rays are abundantly hitting the atmosphere, these missing PeVs gamma sources are very probably mostly absorbed by their photon interactions (photo-pion productions, electron pairs creation) at the source environment and/or along the photon propagation into the cosmic Black Body Radiation (BBR) or into other diffused (radio,infrared,optical) background radiation. Unfortunately PeVs charged cosmic rays, bend and bounded in a random walk by Galactic magnetic fields, loose their original directionality and their astronomical relevance; their resident time in the galaxy is much longer ($\geq 10^3$ - 10^5) than neutral ones, as gamma rays, making the charged cosmic rays more probable to be observed by nearly a comparable ratio. On the contrary astrophysical UHE neutrino signals at 10^{13} eV- 10^{19} eV (or higher GZK energies) are unaffected by any radiation cosmic opacity and may open a very new exciting window to High Energy Astrophysics. Lower energy astrophysical UHE ν at 10^9 eV- 10^{12} eV should also be present, but their signals are (probably) drowned by the dominant diffused atmospheric ν secondaries noises produced by the same charged (and smeared) UHE cosmic rays (while hitting terrestrial atmosphere), the so called atmospheric neutrinos. In a very far corner, at lowest (MeVs) energy windows, the abundant and steady solar neutrino flux and the prompt (but rarer) neutrino burst from a nearby Super-Novae (SN 1987A), have been in last twenty years, already successfully explored.

1.1 The UHE ν Km^3 detectors

The UHE 10^{13} eV- 10^{16} eV ν 's astronomy, being weakly interacting and rarer, may be captured mainly inside huge volumes, bigger than Super-Kamiokande ones; at present most popular detectors consider underground ones (Cubic Kilometer Size like AMANDA-NESTOR)[17] or (at higher

energy $10^{19}\text{eV}-10^{21}\text{eV}$) the widest Terrestrial atmospheric sheet volumes (Auger-Array Telescope or EUSO atmospheric Detectors). Underground km^3 detection is based mainly on ν_μ (above hundred TeVs energies, after their interaction with matter) leading to μ kilometer size lepton tracks [19]. Rarest atmospheric horizontal shower are also expected by ν interactions in air (and, as we shall discuss, in the Earth Crust). While km^3 detectors are optimal for PeVs neutrino muons, the Atmospheric Detectors (AUGER-EUSO like) exhibit a minimal threshold at highest ($\geq 10^{18}\text{eV}$) energies.

1.2 UHECR EeV n and PeV ν Astronomy

Incidentally just around such EeV (10^{18}eV) energies an associated Ultra High Energy Neutron Astronomy might be possible because of the relativistic neutrons boosted lifetime, comparable to our distance from the Galactic center. Therefore UHE neutrons at EeV may be source of part of UHECR data. Indeed a 4% galactic anisotropy and clustering in EeV cosmic rays is recently emerging [1] along our nearby galactic plane. Therefore AGASA might have already experienced a first UHECR-Neutron astronomy **UHENA**. This EeV-UHENA signals must also be source of a parasite ($10^{17} - 10^{16} \text{ eV}$) secondary tails of UHE neutrino $\bar{\nu}_e$ from the same neutron beta decay in flight. Their oscillations in galactic or extragalactic flights (analogous to atmospheric and solar ones) must guarantee the presence of all lepton flavours nearly at equal foot: $\bar{\nu}_e, \bar{\nu}_\mu, \bar{\nu}_\tau$. The latter UHE $\bar{\nu}_\tau$ imprint (added to other local astrophysical UHE ν production) could be already recorded [10] as Upward Tau Air-Showers Terrestrial Gamma Flash (assumed secondaries γ of Upward Tau Air-Showers and Horizontal Tau Air-Showers). These estimates are part of present article below.

1.3 GZK ν Astronomy and Z-Showering

Finally, let us also remind that at highest energy edges ($\geq 10^{19} - 10^{20}\text{eV}$), a somehow correlated New UHE Astronomy is also expected for charged Cosmic Rays; indeed these UHECR have such a large rigidity to avoid any bending by random galactic or extragalactic magnetic fields; being nearly undeflected UHECR should point toward the original sources showing in sky a new astronomical map. (There have been very reiterated attempts to consider extreme galactic or extragalactic coherent magnetic fields able to bend the well known nearby, like Virgo Clusters, M 87 AGN, sources in a diffused map, however their needed magnetic energy density and coherent lengths are in our opinion totally unrealistic). Moreover such UHECR as-

tronomy is bounded by the ubiquitous cosmic $2.75K^o$ BBR screening (the well known Greisen,Zat'sepin,Kuzmin GZK cut-off) limiting its origination inside a very local ($\leq 20Mpc$) cosmic volume. Surprisingly, these UHECR above GZK (already up to day above 60 events) are not pointing toward any known nearby candidate source. Moreover their nearly isotropic arrival distributions underlines and testify a very possible cosmic origination, in disagreement with any local (Galactic plane or Halo) expected footprint by GZK cut-off. A very weak Super-Galactic imprint seem to be present but already above GZK volume. This opened a very hot debate in modern astrophysics known as the GZK paradox. Possible solutions has been found recently beyond Standard Model assuming a non-vanishing neutrino mass. Indeed at such Ultra-High energies, neutrino at ZeV energies ($\geq 10^{21}eVs$) hitting onto relic cosmological light ($0.1 - 4eV$ masses) neutrinos [6] nearly at rest in Dark Hot Halos (galactic or in Local Group) has the unique possibility to produce UHE resonant Z bosons (the so called Z-burst or better Z-Showering scenario): [8], [9], [24],[25]; for a more updated scenario see [14]. Indeed UHE neutrinos are un-effected by magnetic fields and by BBR screening; they may reach us from far cosmic edges with negligible absorption. The UHE Z-shower in its ultra-high energy nucleonic secondary component may be just the observed final UHECR event on Earth. This possibility has been reinforced by very recent correlations (doublets and triplets events) between UHECR directions with brightest Blazars sources at cosmic distances (redshift ≥ 0.1) quite beyond ($\geq 300Mpc$) any allowed GZK cut-off [23] [1],[22]. Therefore there might be a role for GZK neutrino fluxes, either as primary in the Z-Showering scenario or, at least, as (but at lower intensities) necessary secondaries of all those UHECR primary absorbed in cosmic BBR radiation fields. Naturally other solutions as topological defects or primordial relics decay may play a role as a source of UHECR, but the observed clustering [1], [23],[22] seems to favor compact sources. The most recent evidence for self-correlations clustering at 10^{19} , $2 \cdot 10^{19}$, $4 \cdot 10^{19}$ eVs energies observed by AGASA (Teshima, ICRR26 Hamburg presentation 2001) maybe a first reflection of UHECR Z-Showering secondaries: $p, \bar{p}, n, \bar{n}$ [14]. A very recent solution beyond the Standard Model (but within Super-Symmetry) consider Ultra High Energy Gluinos as the neutral particle bearing UHE signals interacting nearly as an hadron in Terrestrial Atmosphere [2]; this solution has a narrow window for gluino masses allowable (and serious problems in production bounds), but it is an alternative that deserves attention. To conclude the puzzle one finally needs to scrutiny the UHE ν astronomy and to test the GZK solution within Z-Showering Models by any independent search on Earth for such UHE neutrinos traces above PeVs reaching either

EeVs-ZeVs extreme energies.

1.4 UHE ν Astronomy by τ Air-Shower

Recently [13],[10] it has been proposed a new competitive (and in our opinion, more convenient) UHE ν detection based on ultra high energy ν_τ interaction in matter and its consequent secondary τ decay in flight while escaping from the rock (Mountain Chains) or water (Sea) in air leading to Upward or Horizontal τ Air-Showers (UPTAU's and HORTAU's),[11],[12]. In a pictorial way one may compare the UPTAU's and HORTAU's as the double bang processes expected in km^3 ice-water volumes [21] : the double bang is due first to the UHE ν_τ interaction in matter and secondly by its consequent τ decay in flight. Here we consider a (hidden) UHE ν -N Bang *in* (the rock-water within a mountain or the Earth Crust) and a τ bang *out* in air, whose shower is better observable at high altitudes. The main power of the UPTAU's and HORTAU's detection is the huge amplification of the UHE neutrino signal, which may deliver almost all its energy in numerous secondaries traces (Cherenkov lights, gamma, X photons, electron pairs, collimated muon bundles). Indeed the multiplicity in τ Air-showers secondary particles, $N_{opt} \simeq 10^{12}(E_\tau/PeV)$, $N_\gamma(< E_\gamma > \sim 10 MeV) \simeq 10^8(E_\tau/PeV)$, $N_{e^-e^+} \simeq 2 \cdot 10^7(E_\tau/PeV)$, $N_\mu \simeq 3 \cdot 10^5(E_\tau/PeV)^{0.85}$ makes easy the UPTAU's-HORTAU's discover. These HORTAU's, also named Skimming neutrinos [15], studied also in peculiar approximation in the frame of AUGER experiment, [4], maybe also originated on front of mountain chains [13], [18] either by $\nu_\tau N$, $\bar{\nu}_\tau N$ interactions as well as by $\bar{\nu}_e e \rightarrow W^- \rightarrow \bar{\nu}_\tau \tau$. This new UHE ν_τ detection is mainly based on the oscillated UHE neutrino ν_τ originated by more common astrophysical ν_μ , secondaries of pion-muon decay at PeVs-EeVs-GZK energies. These oscillations are guaranteed by Super Kamiokande evidences for flavour mixing within GeVs atmospheric neutrino data as well as by most solid and recent evidences of complete solar neutrino mixing observed by SNO detector. HORTAU's from mountain chains must nevertheless occur, even for no flavour mixing, as being inevitable $\bar{\nu}_e$ secondaries of common pion-muon decay chains ($\pi^- \rightarrow \mu^- + \bar{\nu}_\mu \rightarrow e^- + \bar{\nu}_e$) near the astrophysical sources at Pevs energies. They are mostly absorbed by the Earth and are only rarely arising as UPTAU's. Their Glashow resonant interaction allow them to be observed as HORTAU's only within a very narrow and nearby crown edges at horizons (not to be discussed here). At wider energies windows ($10^{14}eV - 10^{20}eV$) only neutrino ν_τ , $\bar{\nu}_\tau$ play a key role in UPTAU's and HORTAU's. These Showers might be easily detectable looking downward the Earth's surface from mountains, planes, balloons or satellites

observer. Here the Earth itself acts as a "big mountain" or a wide beam dump target. The present upward τ at horizons should not be confused with an independent and well known, complementary (but rarer) Horizontal Tau Air-shower originated inside the same terrestrial atmosphere: we shall refer to it as the Atmospheric Horizontal Tau Air-Shower. The same UPTAUS have a less competitive upward showering due to $\nu_e \bar{\nu}_e$ interactions with atmosphere, showering in thin upward air layers [3]: let us label this atmospheric Upward Tau as A-UPTAUs and consider its presence as a very small additional contribute, because rock is more than 3000 times denser than air. Therefore at different heights we need to estimate the UPTAUS and HORTAUs event rate occurring along the thin terrestrial crust below the observer, keeping care of their correlated variables: from a very complex sequence of functions we shall be able to define and evaluate the effective HORTAUs volumes keeping care of the thin shower beaming angle, atmosphere opacity and detector thresholds. At the end of the study, assuming any given neutrino flux, one might be easily able to estimate at each height h_1 the expected event rate and the ideal detector size and sensibility for most detection techniques (Cherenkov, photo-luminescent, gamma rays, X-ray, muon bundles).

2 THE UPTAUS-HORTAUs detection

The τ airshowers are observable at different height h_1 leading to different underneath observable terrestrial areas and crust volumes. HORTAUs in deep valley are also relate to the peculiar geographical morphology and composition [10] and more in detail as discussed below. We remind in this case the very important role of UHE $\bar{\nu}_e e \rightarrow W^- \rightarrow \bar{\nu}_\tau \tau^-$ channels which may be well observable even in absence of any $\nu_\tau, \bar{\nu}_\tau$ UHE sources or any neutrino flavour mixing: its Glashow peak resonance make these neutrinos unable to cross all the Earth across but it may be observable beyond mountain chain [10]; while testing τ air-showers beyond a mountain chain one must keep in mind the possible amplification of the signal because of a possible New TeV Physics (see Fig 5) [10]. In the following we shall consider in general the main $\nu_\tau - N, \bar{\nu}_\tau - N$ nuclear interaction on Earth crust. It should be kept in mind also that UPTAUs and in particular HORTAUS are showering at very low densities and their geometrical opening angle (here assumed at $\theta \sim 1^\circ$) is not in general conical (like down-ward showers) but they are more in a thin fan-like shape (like the observed 8 shaped horizontal Air-Showers). The fan shape is opened by the Terrestrial magnetic field bending. These

UPTAU-SHORTAUs duration time is also much longer than common downward showers because their showering occurs at much lower air density: from micro (UPTAU from mountains) to millisecond (UPTAU and SHORTAUs from satellites) long flashes. Indeed the GRO observed Terrestrial Gamma Flashes, possibly correlated with the UPTAU [10] show the millisecond duration times. In order to estimate the rate and the fluence for of UPTAU and SHORTAUs one has to estimate the observable mass, facing a complex chain of questions, leading for each height h_1 , to an effective observable surface and volume from where UPTAU and SHORTAUs might be originated. From this effective volume it is easy to estimate the observable rates, assuming a given incoming UHE ν flux model for galactic or extragalactic sources. Here we shall only refer to the Masses estimate, unrelated to any UHE ν flux models. These steps are linking simple terrestrial spherical geometry and its different geological composition, high energy neutrino physics and UHE τ interactions, the same UHE τ decay in flight and its air-showering physics at different quota within terrestrial air density. Detector physics threshold and background noises, signal rates have been kept in mind [10], but they will be discussed and explained in forthcoming papers.

3 The Skin Crown Earth Volumes

Let us therefore define, list and estimate below the sequence of the key variables whose dependence (shown below or derived in Appendices) leads to the desired SHORTAUs volumes (useful to estimate the UHE ν prediction rates) summarized in a last Table and in Conclusions. These Masses estimate are somehow an lower bound that ignore additional contribute by more penetrating or regenerated τ . [16]. Let us now show the main functions whose interdependence with the observer altitude lead to estimate the UPTAU and SHORTAUs equivalent detection Surfaces, Volumes and Masses.

1. The horizontal distance d_h at given height h_1 toward the horizons:

$$d_h = \sqrt{(R_{\oplus} + h_1)^2 - (R_{\oplus})^2} = 113 \sqrt{\frac{h_1}{km}} \sqrt{1 + \frac{h_1}{2R_{\oplus}}} \cdot km \quad (1)$$

The corresponding horizontal edge angle θ_h below the horizons ($\pi/2$) is:

$$\theta_h = \arccos \frac{R_{\oplus}}{(R_{\oplus} + h_1)} \simeq 1^o \sqrt{\frac{h_1}{km}} \quad (2)$$




Figure 1: The Upward Tau Air-Shower, UPTAU, and the Horizontal Tau Air-Shower, HORTAU, flashing toward an observer at height h_1 . The HORTAU areas are described for water and rock matter density.

(All the approximations here and below hold for height $h_1 \ll R_\oplus$)

2. The consequent characteristic lepton τ energy E_{τ_h} making decay τ in flight from d_h distance just nearby the source:

$$E_{\tau_h} = \left(\frac{d_h}{c\tau_0} \right) m_\tau c^2 =$$

$$\simeq 2.2 \cdot 10^{18} eV \sqrt{\frac{h_1}{km}} \sqrt{1 + \frac{h_1}{2R}} \quad (3)$$

At low quota ($h_1 \leq$ a few kms) the air depth before the Tau decay necessary to develop a shower correspond to a Shower distance $d_{Sh} \sim 6kms \ll d_h$. More precisely at low quota ($h_1 \ll h_o$, where h_o is the air density decay height= 8.55 km.) one finds:

$$d_{Sh} \simeq 5.96km [1 + \ln \frac{E_\tau}{10^{18}eV}] \cdot e^{\frac{h_1}{h_o}} \quad (4)$$

So we may neglect the distance of the final shower respect to the longest horizons ones. However at high altitude ($h_1 \geq h_o$) this is no longer the case (see Appendix A). Therefore we shall introduce from here and in next steps a small, but important modification , whose physical motivation is just to include the air dilution role at highest quota: $h_1 \rightarrow \frac{h_1}{1+h_1/H_o}$, where , as in Appendix A, $H_o = 23$ km.




Figure 2: The geometrical disposal of the main parameters in the text defining the UPTAU and HORTAU Areas; the distances are exaggerated for simplicity.

Therefore previous definition becomes:

$$E_{\tau_h} \simeq 2.2 \cdot 10^{18} eV \sqrt{\frac{h_1}{1 + h_1/H_o}} \sqrt{1 + \frac{h_1}{2R}} \quad (5)$$

This procedure, applied tacitly everywhere, guarantees that there we may extend our results to those HORTAU at altitudes where the residual air density must exhibit a sufficient slant depth. For instance, highest $\gg 10^{19} eV$ HORTAU will be not easily observable because their τ life distance exceed (usually) the horizons air depth lengths.

3. The parental UHE $\nu_\tau, \bar{\nu}_\tau$ or $\bar{\nu}_e$ energies E_{ν_τ} able to produce such UHE E_τ in matter:

$$E_{\nu_\tau} \simeq 1.2 E_{\tau_h} \simeq 2.64 \cdot 10^{18} eV \cdot \sqrt{\frac{h_1}{km}} \quad (6)$$

4. The neutrino (underground) interaction lengths at the corresponding energies is L_{ν_τ} :

$$\begin{aligned} L_{\nu_\tau} &= \frac{1}{\sigma_{E\nu_\tau} \cdot N_A \cdot \rho_r} \\ &= 2.6 \cdot 10^3 km \cdot \rho_r^{-1} \left(\frac{E_{\nu_h}}{10^8 \cdot GeV} \right)^{-0.363} \end{aligned}$$

$$\simeq 304 km \cdot \left(\frac{\rho_{rock}}{\rho_r} \right) \cdot \left(\frac{h_1}{km} \right)^{-0.1815} \quad (7)$$

For more details see [19], [10]. It should be remind that here we ignore the τ multi-bangs [16] that reduce the primary ν_τ energy and pile up the lower energies HORTAUs (EeV-PeVs).

5. The maximal neutrino depth $h_2(h_1)$ under the chord along the UHE neutrino-tau trajectory of lenght $L_\nu(h_1)$:

$$\begin{aligned} h_2(h_1) &= \frac{L_{\nu_h}^2}{2^2 \cdot 2(R - h_2)} \simeq \frac{L_{\nu_h}^2}{8R_\oplus} \simeq \\ &\simeq 1.81 \cdot km \cdot \left(\frac{h_1}{km} \right)^{-0.363} \cdot \left(\frac{\rho_{rock}}{\rho_r} \right)^2 \end{aligned} \quad (8)$$

See figure 2, for more details. Because the above h_2 depths are in general not too deep respect to the Ocean depths, we shall consider either sea (water) or rock (ground) materials as Crown matter density.

6. The corresponding opening angle observed from height h_1 , δ_{1h} encompassing the underground height h_2 at horizons edge (see Fig.2) and the nearest UHE ν arrival directions δ_1 :

$$\begin{aligned} \delta_{1h}(h_2) &= 2 \arctan \frac{h_2}{2d_h} = \\ &= 2 \arctan \left[\frac{8 \cdot 10^{-3} \cdot \left(\frac{h_1}{km} \right)^{-0.863} \left(\frac{\rho_{rock}}{\rho_r} \right)^2}{\sqrt{1 + \frac{h_1}{2R}}} \right] \\ &\simeq 0.91^\circ \left(\frac{\rho_{rock}}{\rho_r} \right)^2 \cdot \left(\frac{h_1}{km} \right)^{-0.863} \end{aligned} \quad (9)$$

7. The underground chord d_{u_1} (see Fig.2 – 4) where UHE ν_τ propagate and the nearest distance d_1 for τ flight (from the observer toward Earth) along the same d_{u_1} direction, within the angle δ_{1h} defined above, angle below the horizons (within the upward UHE neutrino and HORTAUs propagation line) is:

$$d_{u_1} = 2 \cdot \sqrt{\sin^2(\theta_h + \delta_{1h})(R_\oplus + h_1)^2 - d_h^2} \quad (10)$$

Note that by definition and by construction:

$$L_\nu \equiv d_{u_1} \quad (11)$$




Figure 3: Distances from the observer to the Earth (d_1) for different matter densities or to the Horizons (d_h) for low altitudes

The nearest HORTAUs distance corresponding to this horizontal edges still transparent to UHE τ is:

$$d_1(h_1) = (R_{\oplus} + h_1) \sin(\theta_h + \delta_{1h}) - \frac{1}{2}d_{u_1} \quad (12)$$

Note also that for height $h_1 \geq km$:

$$\begin{aligned} \frac{d_{u_1}}{2} &\simeq (R_{\oplus} + h_1) \sqrt{\delta_{1h} \sin 2\theta_h} \simeq \\ &\simeq 158 \sqrt{\frac{\delta_{1h}}{1^\circ}} \sqrt{\frac{h_1}{km}} km \end{aligned} \quad (13)$$

8. The same distance projected cord $x_1(h_1)$ along the horizontal line (see Fig.2):

$$x_1(h_1) = d_1(h_1) \cos(\theta_h + \delta_{1h}) \quad (14)$$

9. The total terrestrial underneath any observer at height h_1 is A_T :

$$= 2\pi R_{\oplus}^2 (1 - \cos \tilde{\theta}_h) = 2\pi R_{\oplus} h_1 \left(\frac{1}{1 + \frac{h_1}{R_{\oplus}}} \right)$$

$$A_T = 4 \cdot 10^4 km^2 \left(\frac{h_1}{km} \right) \left(\frac{1}{1 + \frac{h_1}{R}} \right) \quad (15)$$

Where $\tilde{\theta}_h$ is the opening angle from the Earth along the observer and the horizontal point whose value is the maximal observable one. At first sight one may be tempted to consider all the Area A_T for UPTAUs and HORTAUs but because of the air opacity (HORTAUs) or for its paucity (UPTAUs) this is incorrect. While for HORTAUs there is a more complex Area estimated above and in the following, for UPTAUs the Area Ring (or Disk) is quite simpler to derive following very similar geometrical variables summarized in Appendix B.

10. The Earth Ring Crown crust area $A_R(h_1)$ delimited by the horizons distance d_h and the nearest distance d_1 still transparent to UHE ν_τ . The ring area $A_R(h_1)$ is computed from the internal angles $\delta\tilde{\theta}_h$ and $\delta\tilde{\theta}_1$ defined at the Earth center (note that $\delta\tilde{\theta}_h = \delta\theta_h$ but in general $\delta\tilde{\theta}_1 \neq \delta\theta_1$).

$$A_R(h_1) = 2\pi R_\oplus^2 (\cos \tilde{\theta}_1 - \cos \tilde{\theta}_h) \quad (16)$$

$$= 2\pi R_\oplus^2 \left(\sqrt{1 - \left(\frac{x_1(h_1)}{R_\oplus} \right)^2} - \frac{R_\oplus}{R_\oplus + h_1} \right) \quad (17)$$

Here $x_1(h_1)$ is the cord defined above.

11. The characteristic interaction lepton tau lenght l_τ defined at the average E_{τ_1} , from interaction in matter (rock or water). These lenghts have been derived by a analytical equations keeping care of the Tau lifetime, the photo-nuclear losses, the electro-weak losses [10]. See figure 5 below.
12. The l_{τ_2} projected along the $\sin(\delta\tilde{\theta}_{h_1})$:

$$\delta\tilde{\theta}_{h_1} \equiv \tilde{\theta}_h - \arcsin \left(\frac{x_1}{R_\oplus} \right) \quad (18)$$

The same quantity in a more direct approximation:

$$\sin \delta\tilde{\theta}_{h_1} \simeq \frac{L_\nu}{2R_\oplus} = \frac{304 km}{2R_\oplus} \left(\frac{\rho_{rock}}{\rho} \right) \frac{h_1}{km}^{-0.1815}.$$

Figure 4: Areas and Angles for UPTAUS-HORTAUS




Figure 5: Lepton τ (and μ) Interaction Lengths for different matter density: R_{τ_0} is the free τ length, $R_{\tau_{New}}$ is the New Physics TeV Gravity interaction range at corresponding densities, $R_{\tau_{Nuc\prime p}}$, [10], see also [5], [7], is the combined τ Ranges keeping care of all known interactions and lifetime and mainly the photo-nuclear one. There are two slightly different split curves (for each density) by two comparable approximations in the interaction laws. R_{Weakp} is the electro-weak Range at corresponding densities (see also [19]); [10].




Figure 6: The $\delta\tilde{\theta}_{h_1}$ opening angle toward Ring Earth Skin for density ρ_{water} and ρ_{rock}

From highest ($h \gg H_o=23\text{km}$) altitude the exact approximation reduces to:

$$\delta\tilde{\theta}_{h_1} \simeq 1^\circ \left(\frac{\rho_{rock}}{\rho} \right) \left(\frac{h_1}{500 \cdot \text{km}} \right)^{-0.1815}$$

Therefore the penetrating τ skin depth $l_{\tau_\downarrow}$ is

$$l_{\tau_\downarrow} = l_\tau \cdot \sin \delta\tilde{\theta}_{h_1} \simeq 0.0462 \cdot l_\tau \left(\frac{\rho_{water}}{\rho} \right) \frac{h_1}{\text{km}}^{-0.1815} \quad (19)$$

Where the τ ranges in matter, l_τ has been calculated and shown in Fig.5.

13. The final analytical expression for the Earth Crust Skin Volumes and Masses under the Earth Skin inspected by HORTAUs are derived combining the above functions on HORTAUs Areas with the previous lepton Tau $l_{\tau_\downarrow}$ vertical depth depths:

$$V_{h_1} = A_R(h_1) \cdot l_{\tau_\downarrow}; \quad (20)$$

$$M_{h_1} = V_{h_1} \cdot \left(\frac{\rho}{\rho_{water}} \right) \quad (21)$$

14. A More approximated but easy to handle expression for Ring area for high altitudes ($h_1 \gg 2\text{km}$ $h_1 \ll R_\oplus$) may be summarized as:

$$A_R(h_1) \simeq 2\pi R_\oplus^2 \sin \theta_h \delta\tilde{\theta}_{h_1} \propto \rho^{-1}$$




Figure 7: Total Area A_T and Ring Areas for two densities A_R at low altitudes




Figure 8: Total Area A_T and Ring Areas for two densities A_R at high altitudes

$$\simeq 2\pi R_{\oplus}^2 \sqrt{\frac{2h_1}{R_{\oplus}}} \left(\frac{\sqrt{1 + \frac{h_1}{2R_{\oplus}}}}{1 + \frac{h_1}{R}} \right) \left(\frac{L_{\nu}}{2R_{\oplus}} \right) \quad (22)$$

At high altitudes the above approximation corrected accordingly to the exact one shown in Figure, becomes:

$$\begin{aligned} A_R(h_1) &\simeq 2\pi R_{\oplus} d_{h1} \delta\tilde{\theta}_{h1} \simeq \\ &\simeq 4.65 \cdot 10^6 \sqrt{\frac{h_1}{500km}} \left(\frac{\rho_{water}}{\rho} \right) km^2 \end{aligned} \quad (23)$$

Within the above approximation the final searched Volume V_{h1} and Mass M_{h1} from where HORTAUs may be generated is:

$$V_{h1} = \frac{\pi}{2} \sqrt{\frac{2h_1}{R_{\oplus}}} \left(\frac{\sqrt{1 + \frac{h_1}{2R}}}{1 + \frac{h_1}{R}} \right) L_{\nu}^2 l_{\tau} \propto \rho^{-3} \quad (24)$$

$$M_{h1} = \frac{\pi}{2} \sqrt{\frac{2h_1}{R_{\oplus}}} \left(\frac{\sqrt{1 + \frac{h_1}{2R}}}{1 + \frac{h_1}{R}} \right) L_{\nu}^2 l_{\tau} \rho \propto \rho^{-2} \quad (25)$$

15. The effective observable Skin Tau Mass $M_{eff.}(h_1)$ within the thin HORTAU or UPTAUs Shower angle beam $\simeq 1^\circ$ is suppressed by the solid angle of view $\frac{\delta\Omega}{\Omega} \simeq 2.5 \cdot 10^{-5}$.

$$\Delta M_{eff.}(h_1) = V_{h1} \cdot \left(\frac{\rho}{\rho_{water}} \right) \frac{\delta\Omega}{\Omega} \quad (26)$$

The Masses $M_{eff.}(h_1)$ for realistic high quota experiment are discussed in the Table below.

4 Summary and Conclusions

The discover of the expected UHE neutrino Astronomy is urgent and just behind the corner. Huge volumes are necessary. Beyond underground km^3 detectors a new generation of UHE neutrino calorimeter lay on front of mountain chains and just underneath our feet: The Earth itself offers huge Crown Volumes as Beam Dump calorimeters observable via upward Tau Air Showers, UPTAU's and HORTAU's. Their effective Volumes as a function of the quota h_1 has been derived by an analytical function variables in equation (17), (20), (23), (28) and Appendix B. These Volumes are discussed below and summarized in the last column of next Table (Fig.9) at the end of the paper. At a few tens meter altitude the UPTAU's and HORTAU's Ring are almost overlapping. At low altitude $h_1 \leq 2km$ the HORTAU's are nearly independent on the ρ matter density: $\Delta M_{eff.}(h_1 = 2km)(\rho_{Water}) = 0.987km^3$ $\Delta M_{eff.}(h_1 = 2km)(\rho_{Rock}) = 0.89km^3$ These volumes are the effective Masses expressed in Water equivalent volumes. On the contrary at higher quotas, like highest Mountain observations sites, Airplanes, Balloons and Satellites, the matter density of the HORTAU's Ring Areas play a more and more dominant role asymptotically *proportional* to ρ^{-2} : $\Delta M_{eff.}(h_1 = 5km)(\rho_{Water}) = 3.64km^3$ $\Delta M_{eff.}(h_1 = 5km)(\rho_{Rock}) = 2.17km^3$ From Air-planes or balloons the effective volumes $M_{eff.}$ increases and the density ρ plays a relevant role. $\Delta M_{eff.}(h_1 = 25km)(\rho_{Water}) = 20.3km^3$ $\Delta M_{eff.}(h_1 = 25km)(\rho_{Rock}) = 6.3km^3$ Finally from satellite altitudes the same effective volumes $M_{eff.}$ are reaching extreme values.

$$\Delta M_{eff.}(h_1 = 500km)(\rho_{Water}) = 150.6km^3$$

$$\Delta M_{eff.}(h_1 = 500km)(\rho_{Rock}) = 30km^3$$

These Masses must be compared with other proposed km^3 detectors, keeping in mind that these HORTAU's signals conserve the original UHE ν direction information within a degree. One has to discriminate HORTAU's (only while observing from satellites) from Horizontal High Altitude Showers (HIAS) [12], due to rare UHECR showering on high atmosphere. One might also remind the UPTAU's (at PeVs energies) volumes as derived in Appendix B and in [10] whose values (assuming an arrival angle $\simeq 45^\circ - 60^\circ$ below the horizons) are nearly *proportional* to the ρ density:

$$\Delta M_{eff.}(h_1 = 500km)(\rho_{Water}) = 5.9km^3$$

$$\Delta M_{eff.}(h_1 = 500km)(\rho_{Rock}) = 15.6km^3$$

These widest Masses values, here estimated analytically for any quota, are offering an optimal opportunity to reveal UHE ν at PeVs and EeVs-GZK energies by crown array detectors (scintillators, Cherenkov, photo-luminescent) facing vertically the Horizontal edges, located at high mountain peaks or at air-plane low sides and finally on balloons and satellites. As it can be seen in last column of final Table , the ratio R between HORTAUs events and Showers over atmospheric UHE ν interaction is a greater and greater number with growing height, implying a dominant role (above two order of magnitude) of HORTAUS grown in Earth Skin Crown over Atmospheric HORTAUs.

5 Event Rate of Upward and Horizontal Tau Air-showers

The event rate for HORTAUs are given by the following expression normalized to any given neutrino flux Φ_ν :

$$\dot{N}_{year} = \Delta M_{eff.} \cdot \Phi_\nu \cdot \dot{N}_o \cdot \frac{\sigma_{E_\nu}}{\sigma_{E_{\nu_o}}} \quad (27)$$

Where the $\dot{N}_o$ is the UHE neutrino rate estimated for km^3 at any given (unitary) energy E_{ν_o} , in absence of any Earth shadow. In our case we shall normalize our estimate at $E_{\nu_o} = 3$ PeVs energy for standard electro-weak charged current in a standard parton model [19] and we shall assume a model-independent neutrino maximal flux Φ_ν at a flat fluence value of nearly $\Phi_{\nu_o} \simeq 3 \cdot 10^3 eV cm^{-2} \cdot s^{-1} \cdot sec^{-1} \cdot sr^{-1}$ corresponding to a characteristic Fermi power law in UHE ν primary production rate decreasing as $\frac{dN_\nu}{dE_\nu} \simeq E_\nu^{-2}$ just below present AMANDA bounds. The consequent rate becomes:

$$\begin{aligned} \dot{N}_{year} &= 29 \frac{\Delta M_{eff.}}{km^3} \cdot \frac{\Phi_\nu}{\Phi_{\nu_o}} \cdot \frac{\sigma_{E_\nu}}{\sigma_{E_{\nu_o}}} \\ &= 29 \cdot \left(\frac{E_\nu}{3 \cdot 10^6 \cdot GeV} \right)^{-0.637} \frac{\Delta M_{eff.}}{km^3} \cdot \frac{\Phi_\nu}{\Phi_{\nu_o}} \end{aligned} \quad (28)$$

For highest satellites and for a characteristic UHE GZK energy fluence $\Phi_{\nu_o} \simeq 310^3 eV cm^{-2} \cdot s^{-1} \cdot sr^{-1}$ (as the needed Z-Showering one), the

consequent event rate observable $\dot{N}_{year}$ above the Sea is :

$$= 12.3 \cdot \left(\frac{E_\nu}{3 \cdot 10^{10} \cdot GeV} \right)^{-0.637} \frac{h}{500km} \cdot \frac{\Phi_\nu}{\Phi_{\nu o}} \quad (29)$$

This event rate is comparable to UPTAUS one and it may be an additional source of Terrestrial Gamma Flashes observed by GRO in last decade [10].

In final Fig. 10 it has been summarized the consequent different detector thresholds (at height $h = 2, 5, 25, 500$ km) assuming a ten years records over different neutrino flux models and in respect with known bounds [20].

6 Appendix A

As soon as the altitude h_1 and the corresponding energy $E_{\tau_{h_1}}$ increases the corresponding air density decreases. At a too high quota there is no more X slant depth for any Air-Showering to develop. Indeed its value is :

$$\begin{aligned} X &= \int_{\frac{d_1}{2} + c\tau\gamma_t}^{d_1 + \frac{d_1}{2}} n_0 e^{-\frac{R_\oplus}{h_0} \left[\sqrt{\left(1 - \frac{h_2}{R_\oplus}\right)^2 + \left(\frac{x}{R_\oplus}\right)^2} - 1 \right]} dx \\ &\simeq \int_{\frac{d_1}{2} + c\tau\gamma_t}^{d_1 + \frac{d_1}{2}} n_0 e^{-\frac{x^2}{2R_\oplus h_0}} dx \leq n_0 h_0 \end{aligned} \quad (30)$$

In order to find this critical height h_1 where the maximal energy HORTAU terminates we remind our recent approximation. The transcendental equation that defines the Tau distance $c\tau$ has been more simplified in:

$$\int_0^{+\infty} n_0 e^{-\frac{\sqrt{(c\tau+x)^2 + R_\oplus^2} - R_\oplus}{h_0}} dx \cong n_0 h_0 A \quad (31)$$

$$\int_0^{+\infty} n_0 e^{-\frac{(c\tau+x)^2}{2h_0 R_\oplus}} dx \cong n_0 h_0 A \quad (32)$$

$$c\tau = \sqrt{2R_\oplus h_0} \sqrt{\ln\left(\frac{R_\oplus}{c\tau}\right) - \ln A} \quad (33)$$

Here $A = A_{Had.}$ or $A = A_\gamma$ are slow logarithmic functions of values near unity; applying known empirical laws to estimate this logarithmic growth (as a function of the X slant depth) we derived respectively for hadronic and gamma UHECR showers [10], [11]:

$$A_{Had.} = 0.792 \left[1 + 0.02523 \ln \left(\frac{E}{10^{19} eV} \right) \right] \quad (34)$$

$$A_\gamma = \left[1 + 0.04343 \ln \left(\frac{E}{10^{19} eV} \right) \right] \quad (35)$$

The solution of the above transcendental equation leads to a characteristic maximal UHE $c\tau_\tau = 546 \text{ km}$ flight distance, corresponding to $E_{1.1} \cdot 10^{19} eV$ energy whose decay occurs at height $H_o = 23 \text{ km}$; from there on the HORTAUS begins to shower. At higher quotas the absence of air density lead to a suppressed development or to a poor particle shower, hard to be detected. At much lower quota the same air opacity filter most of the electromagnetic Shower allowing only to muon bundles to survive at low ($\leq 10^{-3}$) level.

7 Appendix B: The UPTAUS Area

The Upward Tau Air-Showers, mostly at PeV energies, might travel a minimal air depth before reaching the observer in order to amplify its signal. The UPTAUS Disk Area A_U underneath an observer at height h_1 within a opening angle $\tilde{\theta}_2$ from the Earth Center is:

$$A_U = 2\pi R_\oplus^2 (1 - \cos \tilde{\theta}_2) \quad (36)$$

Where the $\sin \tilde{\theta}_2 = (x_2/R_\oplus)$ and x_2 behaves like x_1 defined above for HORTAUS. In general the UPTAUS area are constrained in a narrow Ring (because the mountain presence itself or because the too near observer distances from Earth are encountering a too short air slant depth for showering or a too far and opaque atmosphere for the horizontal UPTAUS):

$$A_U = 2\pi R_\oplus^2 (\cos \tilde{\theta}_3 - \cos \tilde{\theta}_2) \quad (37)$$

An useful Euclidean approximation is:

$$A_U = \pi h_1^2 (\cot \theta_2^2 - \cot \theta_3^2) \quad (38)$$

Where θ_2, θ_3 are the outgoing τ angles on the Earth surface [?].

For UPTAUs (around $3 \cdot 10^{15} eV$ energies) these volumes have been estimated in [?], assuming an arrival values angle $\simeq 45^\circ - 60^\circ$ below the horizons. For two characteristic densities one finds respectively:

$$\Delta M_{eff.}(h_1 = 500km)(\rho_{Water}) = 5.9 \text{ km}^3;$$

$$\Delta M_{eff.}(h_1 = 500km)(\rho_{Rock}) = 15.6 \text{ km}^3$$

Their detection efficiency is displayed in last Figure , and it exceed by more than an order of magnitude, the future ICE-CUBE threshold.

Acknowledgements

The author wishes to thank P.G.De Sanctis Lucentini, C.Leto, M.De Santis for numerical and technical support and Prof.G.Salvini and Prof. B.Mele for useful discussions and comments.

References

- [1] N. Hayashida et al. AGASA collaboration, *Astropart.Phys.* 10 (1999) 303-311.
- [2] Berezhnizky, V.S Gazizov A.Z., Grigorieva S.I., hep-ph/0204357
- [3] Berezhnizky, V.S et al., 1990, *Astrophysics of Cosmic Rays* (North Holland ed.)
- [4] Bertou X., P. Billoir, O. Deligny, C. Lachaud, A. Letessier-Selvon, *Astropart.Phys.* 17 (2002) 183-193
- [5] Becattini, F., Bottai, S. 2001, *Astropart.Phys.*, 15, 323
- [6] Dolgov A. in *Phys.Report.* 2002 (in press); hep-ph/0202122.
- [7] Dutta Iyer, S., Reno, M. H., Sarcevic, I. & Seckel, D. 2001, *Phys.Rev.*, D63, 094020.
- [8] Fargion, D., Salis, A. 1997, *Proc.25th ICRC.* HE-4-6, 153
- [9] Fargion, D., Mele B., Salis, A. 1999, *Ap. J.*, 517, 725; astro-ph/9710029.
- [10] D.Fargion, v570 n2 *ApJ* May 10, 2002; astro-ph/0002453; also astro-ph/9704205;

- [11] D.Fargion, 27th ICRC 2001, HE2.5,p.1297-1300,astro-ph/0106239;
- [12] D.Fargion, 27th ICRC 2001,HE1.8,Germany,903-906,astro-ph/0107094.
- [13] Fargion,D., Aiello, A.,Conversano, R. 1999, 26th ICRC,HE6.1.10, 396-398; astro-ph/9906450.
- [14] Fargion,D., Grossi, M., De Sanctis Lucentini, P.G., Di Troia, C. J.Phys. Soc.Jpn.Vol.70,p.46-57,2001;
- [15] J.L. Feng, Peter Fisher, Frank Wilczek, Terri M. Yu; hep-ph/0105067
- [16] F.Halzen, D. Saltzberg,Phys.Rev.Lett. 81 (1998) 4305-4308
- [17] F. Halzen, astro-ph/0206268
- [18] G.W.S. Hou, M.A. Huang; astro-ph/0204145.
- [19] Gandhi, R., Quigg, C., Reno, M. H., Sarcevic,I. 1998, Phys. Rev. D, 58, 093009
- [20] Kalashev O E, Kuzmin V A, Semikoz D V and Sigl G 2002 Phys. Rev. D 66 063004
- [21] Learned J. G., Pakvasa, S. 1995, Astropart. Phys. 3, 267
- [22] Takeda M. and AGASA Collab.,J.Phys.Soc.Jpn.Vol.70,p.15-21,2001.
- [23] Tinyakov P.,Tkachev I.,J.Phys.Soc.Jpn.Vol.70,p.58-63,2001.
- [24] Yoshida S., Sigl G., Lee S. 1998, Phys.Rev.Lett. 81,5505;
- [25] Weiler T. 1999, Astropart.Phys.11,303.

ρ	h_1	$E_{th}(eV)$	$A_{TOT}(km^2)$	$l_\tau(km)$	$\delta\tilde{\theta}$	$l_\tau \sin \delta\tilde{\theta}(km)$	$A_R(Km^2)$	$\frac{d\Omega}{dV}$	$\Delta V(km^3)$	$V_{eff} = \frac{\Delta V \Delta\Omega}{\Omega}$	$\Delta M(km^3)$	$R = \frac{M_T}{M_{ATM}}$
1	2	$3.12 * 10^{18}$	$8 * 10^4$	21.7	1.31°	0.496	$7.9 * 10^4$	$2.5 * 10^{-5}$	$3.95 * 10^4$	0.987	0.987	49.6
2.65	2	$3.12 * 10^{18}$	$8 * 10^4$	11	0.97°	0.186	$7.2 * 10^4$	$2.5 * 10^{-5}$	$1.34 * 10^4$	0.335	0.89	49.2
1	5	$4.67 * 10^{18}$	$2 * 10^5$	24.3	1.79°	0.76	$1.9 * 10^5$	$2.5 * 10^{-5}$	$1.45 * 10^5$	3.64	3.64	75
2.65	5	$4.67 * 10^{18}$	$2 * 10^5$	12.1	1.07°	0.225	$1.45 * 10^5$	$2.5 * 10^{-5}$	$3.2 * 10^4$	0.82	2.17	59.6
1	25	$8 * 10^{18}$	10^6	27.5	2.36°	1.13	$7.16 * 10^5$	$2.5 * 10^{-5}$	$8.12 * 10^5$	20.3	20.3	113
2.65	25	$8 * 10^{18}$	10^6	13.1	1.08°	0.247	$3.83 * 10^5$	$2.5 * 10^{-5}$	$9.5 * 10^4$	2.4	6.3	65.45
1	500	$1.08 * 10^{19}$	$1.8 * 10^7$	29.4	2.72°	1.399	$4.3 * 10^6$	$2.5 * 10^{-5}$	$6 * 10^6$	150.6	150.6	140
2.65	500	$1.08 * 10^{19}$	$1.8 * 10^7$	13.8	1.07°	0.257	$1.75 * 10^6$	$2.5 * 10^{-5}$	$4.5 * 10^5$	11.3	30	68

Figure 9: The Table of the main parameters leading to the effective HORTAU's Mass from the observer height h_1 , the corresponding τ energy E_τ able to let the τ reach him from the horizons, the Total Area A_{TOT} underneath the observer, the corresponding τ propagation length in matter l_τ , the opening angle toward the Crown from the Earth $\delta\tilde{\theta}_{h_1}$ and l_τ just orthogonal in the matter $l_{\tau_\perp} = l_\tau \cdot \sin \delta\tilde{\theta}_{h_1}$, the Ring Areas for two densities A_R at characteristic high altitudes h_1 , the corresponding effective Volume V_{eff} . and the consequent Mass ΔM_{eff} . (within the narrow τ Air-Shower solid angle) as a function of density ρ and height h_1 . In the last Column the Ratio $R = M_T/M_{ATM}$ define the ratio of HORTAU's produced within the Earth Crown Skin over the atmospheric ones: this ratio nearly reflects the matter over air density and it reaches nearly two order of magnitude.




Figure 10: UPTAUS (lower bound on the center) and HORTAUS (right parabolic curves) sensibility at different observer heights h (2, 5, 25, 500km) assuming a km^3 scale volume (see Table above) adapted over a present neutrino flux estimate in Z-Shower model scenario [20], [9] for light ($0.4 - 0.04$ eV) neutrino masses m_ν ; two corresponding density contrast has been assumed [14]; the lower parabolic bound thresholds are at different operation height, in Horizontal (Crown) Detector facing toward most distant horizons edge; these limits are fine tuned (as discussed in the text); we are assuming a duration of data records of a decade comparable to the BATSE record data (a decade). The paraboloid bounds on the EeV energy range in the right sides are nearly un-screened by the Earth opacity while the corresponding UPTAUS bounds in the center below suffer both of Earth opacity as well as of a consequent shorter Tau interaction length in Earth Crust, that has been taken into account. [10], .